



## Soils Formed In Marine Sediments Containing Glauconite

Series	Solum (in)	Parent Material
<b>Well drained low glauconite</b>		
Colts Neck	30-50	Loamy marine sediments that contain glauconite
Freehold	30-45	Sandy marine sediments which have small to moderate amounts of glauconite
Howell	40-60	Marine sediments with low glauconite
Swedesboro	30-45	Loamy and sandy marine sediments that contain glauconite
<b>Moderately well drained low glauconite</b>		
Holmdel	30-45	Loamy glauconite bearing marine and/or fluviomarine sediments
Sharptown	20-48	Silty eolian deposits underlain by loamy marine sediments that contain glauconite
Woolwich	30-45	Sandy marine sediments that contain glauconite
<b>Well drained moderate glauconite</b>		
Collington	28-44	Glauconite bearing eolian and/or fluviomarine deposits
Tinton	30-50	Sandy eolian deposits over glauconite bearing fluviomarine deposits
<b>Moderately well drained moderate glauconite</b>		
Adelphia	25-60	Glauconite bearing eolian and/or fluviomarine deposits
Donlonton	40-60	Old marine sediments containing moderate amounts of glauconite
Pemberton	24-48	Sandy eolian deposits underlain by loamy alluvium derived from marine sediments that contain glauconite
<b>Poorly drained moderate glauconite</b>		
Shrewsbury	24-40	Marine sediments containing moderate amounts of glauconite
<b>Very poorly drained with moderate glauconite</b>		
Keansburg	24-40	Water laid deposits containing low to moderate amounts of glauconite
<b>Well drained with high or very high glauconite</b>		
Monmouth	35-55	Fine-textured marine sediments high in glauconite
<b>Moderately well drained with high or very high glauconite</b>		
Marlton	30-48	Clayey glauconite marine and/or fluviomarine deposits
<b>Somewhat poorly drained with high or very high glauconite</b>		
Kresson	24-40	Clayey glauconite marine and/or fluviomarine deposits
<b>Poorly drained with high or very high glauconite</b>		
Colemantown	26-40	Glauconite bearing fluviomarine deposits